

# Accepted Manuscript

Conservative management of temporomandibular dysfunction: A literature review with implications for clinical practice guidelines

Raymond Butts, DPT, PhD, MSc, Faculty, Senior Instructor, James Dunning, DPT, MSc, FAAOMPT, Director, President, Ron Pavkovich, DPT, OCS, FAAOMPT, Faculty, Senior Instructor, Jersey Mettill, BS, Medical Student, Firas Mourad, DPT, OCS, FAAOMPT, Faculty, Senior Instructor



PII: S1360-8592(17)30121-3

DOI: [10.1016/j.jbmt.2017.05.021](https://doi.org/10.1016/j.jbmt.2017.05.021)

Reference: YJBMT 1544

To appear in: *Journal of Bodywork & Movement Therapies*

Received Date: 19 January 2017

Revised Date: 22 May 2017

Accepted Date: 26 May 2017

Please cite this article as: Butts, R., Dunning, J., Pavkovich, R., Mettill, J., Mourad, F., Conservative management of temporomandibular dysfunction: A literature review with implications for clinical practice guidelines, *Journal of Bodywork & Movement Therapies* (2017), doi: 10.1016/j.jbmt.2017.05.021.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Conservative Management of Temporomandibular Dysfunction: A Literature  
Review with Implications for Clinical Practice Guidelines**

**PART 2**

**Authors:**

**Raymond Butts**, DPT, PhD, MSc, Faculty, American Academy Of Manipulative Therapy Fellowship in Orthopedic Manual Physical Therapy. Senior Instructor, Spinal Manipulation Institute & Dry Needling Institute, Montgomery, Alabama

**James Dunning**, DPT, MSc, FAAOMPT, Director, American Academy Of Manipulative Therapy Fellowship in Orthopedic Manual Physical Therapy. President, Spinal Manipulation Institute & Dry Needling Institute, Montgomery, Alabama

**Ron Pavkovich**, DPT, OCS, FAAOMPT, Faculty, American Academy Of Manipulative Therapy Fellowship in Orthopedic Manual Physical Therapy. Senior Instructor, Spinal Manipulation Institute & Dry Needling Institute, Montgomery, Alabama

**Jersey Mettelle**, BS, Medical Student, Philadelphia College of Osteopathic Medicine, Suwanee, Georgia

**Firas Mourad**, DPT, OCS, FAAOMPT, Faculty, American Academy Of Manipulative Therapy Fellowship in Orthopedic Manual Physical Therapy. Senior Instructor, Spinal Manipulation Institute & Dry Needling Institute, Montgomery, Alabama

**Corresponding Author:**

Dr. Raymond Butts

American Academy of Manipulative Therapy

1036 Old Breckenridge, Ln

Montgomery, Alabama, United States Of America

**Postal Code:** 36117

**Tel:** 803-422-3954

**Email:** [buttsraymond@yahoo.com](mailto:buttsraymond@yahoo.com)

**Abstract**

The effective management of temporomandibular dysfunction (TMD) requires a thorough understanding of the pathoanatomic factors that drive the underlying condition. After reviewing the etiology associated with TMD in Part 1 of this narrative review, the temporomandibular joint capsule, articular disc and muscles of mastication emerged as key players. Part 2 focuses on conservative treatment strategies best able to reduce the pain and disability associated with TMD. A review of the literature revealed limited support of strengthening exercises targeting the muscles of mastication. There was also limited evidence for manual soft tissue work targeting muscles of mastication, which may be specifically related to the limited accessibility of the pterygoid muscles to palpation. For the reduction of pain, there was little to no evidence supporting splint therapy and electrophysical modalities, including laser therapy, ultrasound, TENs and iontophoresis. However, for the reduction of pain and disability, non-thrust mobilization and high-velocity, low amplitude thrust manipulation techniques to the TMJ and/or upper cervical articulations that directly and indirectly target the TMJ joint capsule were generally supported in the literature. Studies that used dry needling or acupuncture of the lateral pterygoid and posterior peri-articular connective tissue also led to significant improvements in pain and disability in patients with TMD. Thus, the most effective conservative management of TMD seems to be techniques best able to impact anatomic structures directly related to

the etiology of TMD, to include the joint capsule, articular disc and muscles of mastication, specifically the superior and inferior head of the lateral pterygoid.

### **Introduction**

According to Shaffer et al., physical therapy is the preferred conservative approach for treating TMD, (Shaffer et al., 2014) as it facilitates multi-modal treatment that addresses patient specific impairments (Guarda-Nardini et al., 2016). However, successful management requires treatment of anatomical structures that are consistent with the underlying condition. In this context, the muscles of mastication, (Ariji et al., 2015; Hiraba et al., 2000; Murray & Peck, 2007; Murray et al., 2004; Peck et al., 2008; Pihut et al., 2016) joint capsule (Friedman, 1997; Mapelli et al., 2016; Saghafi & Curl, 1995) and cervical spine (Fernandez-de-Las-Penas et al., 2010; Guarda-Nardini et al., 2016; Jayaseelan & Tow, 2016; Mansilla-Ferragut et al., 2009) are high-value targets that likely require further consideration.

### **Temporomandibular Dysfunction and Exercise**

Given the reciprocal actions of the superior and inferior head of the lateral pterygoid, (Desmons et al., 2007; Hiraba et al., 2000; Mahan et al., 1983; McNamara, 1973; Murray et al., 2004) it is plausible that decreased activity of the superior head agonist results in increased activity in the inferior head antagonist, a theory consistent with pain adaptation. Similarly, the decreased activity of the superior head could result in overuse of the inferior head, resulting in an energy deficit and setting the stage for muscle hypertonicity via the Vicious Cycle Model. Notably, both models result in overactive muscles of mastication, a problem further propagated by psychosocial

factors (Peck et al., 2008). Perhaps this explains the limited evidence for strengthening muscles of mastication to treat TMD. That is, the purposeful activation of muscles that are already overactive (Lauriti et al., 2014; Tosato Jde et al., 2015) may not be advantageous and does not seem to be strongly supported by the literature (Mulet et al., 2007). Notably, a recent RCT by Bae et al. found masticatory relaxation exercises to be more effective for decreasing the pain and limited ROM associated with TMD than active exercises (Bae & Park, 2013).

The Rocabado exercise routine, (Rocabado & Iglarsh, 1991) which addresses the rest position of the tongue, control of TMJ rotation during mouth opening, rhythmic stabilization of the mandible, head flexion stabilization, lower cervical retraction and shoulder girdle retraction is the most widely used exercise routine used by physical therapists to treat TMD (Shaffer et al., 2014). However, the only study that has examined its effectiveness concluded that that the exercise routine provided no additional therapeutic benefit to the rehabilitation process (Mulet et al., 2007; Shaffer et al., 2014). Kraus (Kraus, 1988, 2004) also proposed a 3-prong approach to TMD, including inhibition of excessive activity of muscles of mastication, mandibular neuromuscular control and isometric exercises to counter joint clicking, muscle asymmetry and spasms, but the regiment has never been studied or validated (Shaffer et al., 2014). According to a systematic review by Shaffer et al., ‘...no evidence exists to direct clinicians toward which exercises, if any, may be useful in the conservative management of TMD. Additionally, because many, if not most, patients with TMD over-recruit their muscles of mastication, it may be more advantageous to

focus on relaxation techniques and patient education than therapeutic exercise' (Shaffer et al., 2014).

In a systematic review of 7 studies that investigated therapeutic exercises for TMD, which included stretching, relaxation, coordination, strengthening and endurance, Moraes et al. found exercises to be effective for the treatment of 'muscular TMD' (Moraes, 2013). However, the authors noted significant issues with internal validity and an over reliance on clinical experience (Moraes, 2013). Notably, all 7 studies included by Moraes et al. (Moraes, 2013) combined therapeutic exercise with other conservative procedures, making it difficult, if not impossible, to draw any firm conclusions about the usefulness of exercise for TMD. Therefore, it is not surprising that the most recent systematic review on TMD by Armijo-Olivo et al. reported 'great uncertainty' for the effectiveness of exercises in treating TMD (Armijo-Olivo et al., 2016). Although the investigators found 'promising' effects for manual therapy and manual therapy combined with exercise, the use of exercise alone was not superior to other conservative treatments for TMD (Armijo-Olivo et al., 2016). Perhaps the masticatory muscle weakness identified by many physical therapists is more likely the product of pain inhibition instead of true weakness. Given that both TMD pain and experimental jaw-muscle pain have been shown to result in decreased EMG activity during clenching, pain inhibition seems to play a role (Castroflorio et al., 2012; Pinho et al., 2000). As such, perhaps physical therapy outcomes would improve if more attention were placed on the pain associated with TMD instead of muscle weakness, the product of pain. (Bae & Park, 2013; Moller et al., 1984)

### **Temporomandibular Dysfunction and Soft Tissue Release**

A systematic review by Turp & Minagi reported a lack of validity and poor inter-examiner reliability for manual palpation of the lateral pterygoid muscle intra-orally and recommended that the procedure be abandoned (Turp & Minagi, 2001). However, with the exception of one lateral head radiograph on a single subject, (Johnstone & Templeton, 1980) the 5 RCTs included by Turp & Minagi used cadavers, and the results may not be generalizable to live patients. Notably, a recent study by Stelzenmueller et al. found that palpation of the lateral pterygoid is 'basically feasible' in both live patients and cadavers with 'exact knowledge of muscle topography and the intraoral palpation pathway' (Stelzenmueller et al., 2016). However, the reliability of clinicians to follow the complex series of instructions required to negotiate the intraoral palpation pathway and bypass the medial pterygoid is unknown (Stelzenmueller et al., 2016). Moreover, while Stelzenmueller and colleagues demonstrate an ability to contact the lateral pterygoid with the tip of either digit three or five, (Stelzenmueller et al., 2016) it is doubtful whether such limited contact would translate into meaningful clinical outcomes. Given the lack of inter-examiner reliability of localizing trigger points in the upper trapezius, (Sciotti et al., 2001) a fully accessible muscle, it also seems highly unlikely that clinicians are able to reliably locate trigger points in the lateral pterygoid and needle them consistently, as a number of authors have claimed (Gonzalez-Perez et al., 2012; Gonzalez-Perez et al., 2015).



The inaccessibility of the lateral pterygoid to palpation may also account for the limited evidence in the treatment of TMD with soft tissue mobilization (Miernik et al., 2012). The most recent systematic review and meta-analysis by Calixtre et al. reported a limited number of studies and low to moderate evidence that myofascial release and massage are better than control and moderate evidence that they are as effective as Botox injections in the masseter and temporalis muscle (Calixtre et al., 2015; Guarda-Nardini et al., 2012). However, given that a clinically relevant level of pain in the masseter muscles has only a minor impact in the performance of the masseter and temporalis, (Manfredini et al., 2013; Shimada et al., 2015) manual soft tissue techniques targeting those muscles may not be useful for more than symptom relief. That is, manual soft tissue techniques may have a positive effect on more accessible muscles such as the temporalis and masseter, thereby improving symptoms associated with TMD, but they are unlikely to reach the pterygoid muscles that are not only more intimately related with the joint itself, but also more likely one of the primarily etiologic factors in TMD (Gauer & Semidey, 2015; Lafreniere et al., 1997; Liu et al., 1989; Scully, 2008, 2013).

### **Temporomandibular Dysfunction and Electrophysical Modalities**

Interestingly, McNeely et al. (2006) found no evidence to support the use of any electrophysical modalities commonly used by physical therapists such as pulsed radio frequency energy, biofeedback, laser therapy and TENS to reduce pain associated with TMD (McNeely et al., 2006). Since the ultrasound penetration half depth into a perpendicular muscle at 1 MHz and 3 MHz is 0.9 and 0.3 cm, respectively, (Cameron,

2003) and laser intensity is reduced by 90% at a tissue depth of 1cm, (Ayyildiz et al., 2015) very little of the sound waves or energy likely reaches the TMJ or pterygoids. Notably, a number of systematic reviews have reported little to no evidence to support using ultrasound for musculoskeletal conditions, (Robertson & Baker, 2001; Shanks et al., 2010; van der Windt et al., 1999) in general, to include TMD (Shaffer et al., 2014). Also, while the depth of penetration of TENs depends on the size and distance between surface electrodes, (Cameron, 2003) Fuglevand et al. reported that only motor units of muscle fibers within 1-1.2cm from the surface electrodes are influenced by the modality (Fuglevand et al., 1992). Thus, much like soft tissue techniques that have a limited ability to reach the pterygoids and TMJ capsule, the source of the problems according to Scully's model of TMD, (Scully, 2008, 2013) electrophysical modalities likely provide very limited relief to patients suffering from TMD.

A number of studies have also considered the use of iontophoresis for treating TMD (Braun, 1987; Furto et al., 2006; Mina et al., 2011; Reid et al., 1994; Schiffman et al., 1996). While dexamethasone can be delivered to depths of up to 17 mm, (Anderson et al., 2003a; Anderson et al., 2003b) the absorption of the drug into target tissue is variable (Smutok et al., 2002). Clinically, studies that have used iontophoresis for TMD (Furto et al., 2006; Mina et al., 2011; Reid et al., 1994; Schiffman et al., 1996) report modest improvements in function but not pain (Shaffer et al., 2014; Wieckiewicz et al., 2015). In a case series of patients with TMD, Furto et al. used a multimodal approach that included iontophoresis with dexamethasone and concluded, patients with TMD who are treated with a rehabilitation program

including manual physical therapy interventions plus exercise, with or without iontophoresis with dexamethasone, can demonstrate clinically meaningful improvements in disability and overall perceived change in a relatively short period of time' (Furto et al., 2006). Given that Reid et al. found iontophoresis to be no better than saline for the treatment of TMD, (Reid et al., 1994) more evidence is required to fully justify the use of the modality, clinically.

### **Temporomandibular Dysfunction and Splint Therapy**

In a 2004 Cochrane review, Al-Ani et al. found insufficient evidence to support the use of splint therapy for the treatment of TMD (Al-Ani et al., 2004). After treating 80 consecutive patients with TMD, Niemela reported that splint treatment, counseling and masticatory muscle exercises were not more beneficial than counseling and masticatory muscle exercises alone. Similarly Nagata et al. found no added short-term benefit of splint therapy in TMD patients receiving multi-modal therapy, including self-exercise, cognitive therapy, self-management education and manipulation (Nagata et al., 2015). Quintus et al. further evaluated the long-term effect of splint therapy. After 1 year, 27.6% of TMD patients that received splint treatment and 37.5% of TMD patients that received counseling and instructions for increasing masticatory muscles exercises, respectively, reported 'very good' treatment effects (Quintus et al., 2015). Even though 16/40 patients in the counseling and exercise group were switched to the splint therapy group because of painful symptoms associated with TMD, both groups experienced only a modest reduction in

pain. Moreover, splint therapy did not outperform counseling and instructions for self-exercise (Qvintus et al., 2015).

### **Temporomandibular Dysfunction and Joint Mobilization**

Non-thrust, joint mobilization has been found to improve the extensibility of noncontractile tissue and increase range of motion while decreasing pain and disability via peripheral, spinal and supraspinal mechanisms (Bialosky et al., 2009; Shaffer et al., 2014). However, very few studies presently exist that have investigated the use of mobilization, independent of other conservative treatments, for TMD. Therefore, after reviewing systematic reviews, RCTs and clinical guidelines related to the use of mobilization for various musculoskeletal disorders, Bronfort et al. reported inconclusive evidence for the use of joint mobilization for TMD (Bronfort et al., 2010). Nevertheless, Taylor et al. used EMG to demonstrate a significant decrease in masseter activity and improved mandibular ROM following passive mobilization compared to sham (Taylor et al., 1994). Given the more intimate relationship of the superior and inferior head of lateral pterygoid to the TMJ capsule, joint mobilization may have an even stronger impact on these muscles.

The use of joint mobilization in combination with other conservative treatment has been evaluated in a number of studies on patients with TMD, and they have, almost without exception, reported significant improvements in pain and mandibular range of motion. In one study, Carmeli et al. treated 36 patients with TMD with either splint therapy or TMJ mobilization and exercise (Carmeli et al., 2001). Joint mobilizations

were progressively advanced from Grade I to Grade IV over 15 treatments, while exercises focused on active mouth opening, closing and lateral deviation X10 repetitions, 6 times per day. After 4 weeks of treatment, only the group receiving mobilization and exercise reported a significant between group improvement in pain and disability (Carmeli et al., 2001).

### **Temporomandibular Dysfunction and Joint Manipulation**

High-velocity low-amplitude (HVLA) thrust manipulation has been shown to increase afferent discharge rates of mechanoreceptors to quiet alpha motor neuron pool activity at the level of the spinal cord and subsequently decrease muscle activation levels (Colloca et al., 2004). A number of case studies have successfully used HVLA thrust manipulation to treat patients with TMD, reporting improvements in pain and disability (Ogawa et al., 2015; Rubis et al., 2014; Yabe et al., 2014). While Gotou et al. reported a significant improvement in condylar movement using JOG-HVLA thrust manipulation (i.e. combination of pivot closing, side-to-side and opening HVLA thrust techniques) to treat 34 patients with TMD and limited mouth opening (<35mm), (Gotou et al., 2010) the most recent systematic review of only 2 medium quality studies on HVLA thrust manipulation of the TMJ by Alves et al. found insufficient evidence to support using the technique for patients with TMD (Alves et al., 2013).

Interestingly, there is considerably more evidence for manipulating the cervical spine (i.e. HVLA thrust manipulation) to treat symptoms associated with TMD.

Anatomically, the spinal aspect of the trigeminocervical nucleus overlaps with the dorsal horns in the upper cervical spine (Chua et al., 2012; Lin, 2014; Spadaro et al., 2014). Thus, nociceptive afferents from the anatomical structures in the upper neck can often be confused with trigeminal pain, to include the TMJ (Chua et al., 2012; Lin, 2014; Spadaro et al., 2014). Notably, Fernandez de-las-penas et al. reported a significant correlation in 25 patients with TMD and myofascial trigger points in the muscles of the neck and shoulder regions (Fernandez-de-Las-Penas et al., 2010). As such, manual techniques such as HVLA thrust manipulation that target the upper cervical spine region (i.e. C0-1, and C1-2) may also effectively inhibit pain and quiet hypertonic muscles of mastication, to include the pterygoids.

A recent systematic review by Calixtre et al. reported low to high evidence for the use of non-thrust mobilization or HVLA thrust manipulation of the cervical spine compared to control in the treatment of TMD (Calixtre et al., 2015). Monaco et al. found a significant increase in mouth opening and opening velocity per kinesiographic data following osteopathic manipulation in 28 children with TMD (Monaco et al., 2008). Moreover, Oliveira-Campelo et al. treated 121 volunteers with latent trigger points in the masseter and reported that a single session of either occipito-atlantal (C0-1) HVLA thrust manipulation or soft tissue release of the suboccipital muscles resulted in an immediate increase in pressure pain thresholds over the masseter and an improvement in mouth opening compared to control (Oliveira-Campelo et al., 2010). Mansilla-Farragut et al. also treated 37 women with chronic neck pain with a single session of bilateral OA HVLA thrust manipulation and

demonstrated an immediate improvement in mouth opening and reduction in pressure pain thresholds over the sphenoid bone compared to control (Mansilla-Ferragut et al., 2009).

Taken together, these studies seem to demonstrate a clear connection between neck pain and the trigeminal nerve, which provides sensation to face, and specifically to the TMJ via the auriculotemporal branch of the mandibular branch of the trigeminal nerve and motor innervation to the muscles of mastication (Fernandez-de-Las-Penas et al., 2010; Moore & Dalley, 2006). While spinal manipulation targeting the upper cervical spine has been shown to improve pain and motor performance in patients with TMD, the longevity of the changes is presently unknown. However, given that Dunning et al. reported a significant improvement in neck pain, disability and motor performance of the deep cervical neck flexors 48-hours following spinal manipulation at C1-C2 and T1-T2 compared to grade IV mobilization, the changes are likely not transient (Dunning et al., 2012).

### **Temporomandibular Dysfunction and Joint Mobilization and/or Manipulation**

A recent systematic review by Martins et al. notes a general consensus in the literature and B-level of evidence supporting TMJ mobilization (or manipulation), a manual technique applied to the 'craniocervical mandibular system' and commonly combined with 'multi-modal therapy', for improving pain and ROM in patients with TMD (Martins et al., 2016). Moreover, the authors reported a large effect and a significant difference in active mouth opening and pain reduction of manual therapy compared to other active interventions, including home exercise, massage, usual care,

sham treatment and splint therapy (Martins et al., 2016). Interestingly, a systematic review and meta-analysis by Armijo-Olivo et al. (2016) reported that mandibular, postural and cervical exercises were superior to other active conservative treatments for TMD (Armijo-Olivo et al., 2016). However, manual therapy, including manipulation and/or mobilization to the orofacial and C-spine, was effective when administered alone or in combination with exercise (Armijo-Olivo et al., 2016). In general, a 5mm improvement in mouth opening is considered a clinically relevant change; (Kropmans et al., 1999a; Kropmans et al., 1999b) while postural exercises and either jaw or jaw and cervical exercises reported a 5.54 and 5.94mm improvement, respectively, manual therapy led to a 17.33mm improvement (Armijo-Olivo et al., 2016; Kropmans et al., 1999a; Kropmans et al., 1999b). That is, the addition of exercises to manual therapy for the treatment of TMD did not seem to provide an added clinical advantage, suggesting that mobilization (or manipulation) may provide a key component of the intervention (Armijo-Olivo et al., 2016).

### **Temporomandibular Dysfunction and Needling Modalities**

While the terminology, philosophy and theoretical constructs differ between Western-based DN and traditional Chinese acupuncture, the procedure of inserting monofilament needles is similar (Dunning et al., 2014). Therefore, manual and electric DN are used synonymously with manual and electroacupuncture, respectively, throughout this section to describe needling procedures that penetrate the skin in the absence of medicine or injectate (Butts et al., 2016; Dunning et al., 2014).



While acupuncture specifically facilitates direct contact with the TMJ joint capsule and the superior and inferior heads of the lateral pterygoid muscle, many of the controlled trials that have attempted to treat TMD with acupuncture have reported modest outcomes (McNeely et al., 2006). A systematic review by List and Axelson did not support acupuncture for TMD more than occlusal appliances, behavioral therapy, exercise, posture training and pharmaceutical meds (List & Axelsson, 2010). Similarly, Jung et al. published a systematic review of 7 randomized control trials and found only limited evidence for the use of acupuncture for the symptomatic treatment of TMD (Jung et al., 2011). However, methodological issues stand out in the Jung et al., systematic review (Jung et al., 2011). First, a penetrating placebo was used in one of the 7 included, and this comparison treatment was likely not inert (Dincer & Linde, 2003). Second, only 1 of the 7 included trials considered needle manipulation, which is likely an important omission considering the soft-tissue (i.e. mechanotransduction) (Langevin et al., 2006; Langevin et al., 2005; Langevin et al., 2011) vascular, (Sandberg et al., 2003; Yim et al., 2007) peripheral analgesic (Choi et al., 2013; Goldman et al., 2010; Takano et al., 2012) and central descending pain inhibitory effects (Lundeberg & Lund, 2007) that have been linked with needle manipulation (Kong et al., 2007; Zhou & Benharash, 2014). Third, in 4 of the trials, the longest outcome was following just one acupuncture session—likely an under dosing of the intervention (White et al., 2008; White et al., 2007). Fourth, and perhaps most surprising, was the fact that 60 of the 91 points needled across the 7 trials included in Jung et al. were in distal points of the hand (LI4, SI3 and SI2) instead

of the muscles of mastication or the peri-articular connective tissue of the TMJ. In fact, only 13 of the 91 points were consistent with the masseter (ST6) and 6 of the 91 points were consistent with masseter and lateral pterygoid (ST7) (Jung et al., 2011). Similarly, a more recent study by Grillo et al. compared acupuncture with a flat occlusal plane appliance in 40 women with TMD and reported no between-group difference (Grillo et al., 2015). Yet, only 1 out of 10 acupoints used (i.e. SI19) was associated with the TMJ, and none penetrated muscles of mastication (Grillo et al., 2015).

Notably, La Touche et al. reported that local acupoints (i.e. acupoints anatomically positioned near the symptomatic TMJ) achieved the greatest pain reduction (La Touche et al., 2010). More specifically, 2 of the 4 randomized control trials included in the La Touche et al. systematic review produced both statistically significant and clinically meaningful changes in pain associated with TMD using acupoints ST6 (i.e. the lower, posterior border of the masseter muscle) and ST7 (i.e. the mandibular notch, one index finger forward of the tragus, within the deep masseter when inserted superficially and/or the lateral pterygoid muscle when inserted deeply) (La Touche et al., 2010). (See Figure 1)

Cho and Whang published a systematic review of 15 randomized control trials of 808 patients with TMD; (Cho & Whang, 2010) however, in order to combat heterogeneity of various interventions, they qualitatively grouped patients instead of statistically pooling the data. Unlike Jung et al., (Jung et al., 2011) 2 of the 3 most common

acupoints used among the 14 trials were points consistent with the masseter and lateral pterygoid (ST6 and ST7). Notably, the authors concluded that there is moderate evidence that acupuncture is superior to non-penetrating placebo, wait list control, pharmaceutical treatment and physical therapy, to include ultrasound, massage and manual therapy, for the treatment of TMD. In a more recent systematic review of 21 studies, Porporatti et al. also reported significant improvements in pain, joint movement and oral function along with decreased hypertonicity of masticatory muscles secondary to acupuncture (Porporatti et al., 2015). As in Cho and Whang, Porporatti et al. found that ST6 and ST7 were 2 of the 3 most commonly used acupoints for TMD. In addition, the authors suggested that the temporalis muscle also be needled at Taiyang (one index finger posterior of the lateral orbit of the eye) (Porporatti et al., 2015). Given that ST5 (lower, anterior border of the masseter) and ST8 (anterosuperior aspect of the temporalis) correspond to the masseter and temporalis muscle, respectively, (Deadman et al., 2007) these acupoints may also be appropriate needle locations for patients with TMD. (See Figure 1)

While myofascial trigger points in the muscles of mastication can be a source of pain in TMD, (Diracoglu et al., 2012; Gonzalez-Perez et al., 2012; Gonzalez-Perez et al., 2015) there is a lack of robust, empirical evidence validating the clinical diagnostic criteria for identifying trigger points, as originally proposed by Travell and Simons (Simons et al., 1999; Simons & Travell, 1983) and Fischer (Fischer, 1988; Tough et al., 2007) (Sciotti et al., 2001). Nevertheless, the goal of needling trigger points associated with TMD requires clinicians to focus their needles on the key muscles of

mastication (Diracoglu et al., 2012; Gonzalez-Perez et al., 2012; Gonzalez-Perez et al., 2015). As a result, the limited number of studies that have investigated the use of myofascial trigger point (MTrP) dry needling to treat TMD have reported, in the main, positive outcomes in terms of statistically significant and clinically meaningful between-group reductions in both pain and disability.

Interestingly, McMillan et al. found that MTrP dry needling and simulated procaine, simulated MTrP dry needling and procaine, and simulated MTrP dry needling and procaine targeting trigger points in the masseter muscle resulted in modest and statistically equivalent changes in pain intensity and unpleasantness scores (McMillan et al., 1997). In contrast, Fernandez-Carnero et al. found a significant increase in pressure pain thresholds and maximal jaw opening compared to sham after needling trigger points in the masseter muscle (Fernandez-Carnero et al., 2010). Given that the authors performed deep dry needling into the masseter at a depth consistent with the trigger point, it would be interesting to know how often the medial and lateral pterygoid muscles were penetrated. Gonzalez-Perez et al. also performed MTrP dry needling once per week for 3-weeks on muscles of mastication with palpable trigger points in 36 patients with TMD but specifically targeted the inferior head of the lateral pterygoid at an insertion site consistent with the ST7 acupoint (Mesa-Jimenez et al., 2015). While the study did not include a control group, the authors reported a statistically significant within-group improvement in pain and jaw movement compared to baseline directly following treatment and at 6 months post treatment. In a more recent study by Gonzalez-Perez et al., the authors reported

significant between-group reductions in pain and improvements in jaw movements that lasted up to 70 days following MTrP dry needling targeting the lateral pterygoid compared to control (methocarbamol/paracetamol every 6-hours for 3-weeks) (Gonzalez-Perez et al., 2015). Collectively, these findings suggest that the masseter and temporalis play significant roles in TMD; however, and in large part, the primary etiology and the treatment focus for the pain and disability associated with TMD is likely the lateral pterygoid muscle (Scully, 2008, 2013).

While no studies have directly investigated the use of dry needling or acupuncture to improve the health of intra- and peri-articular aspects of the TMJ itself in patients with TMD, there is evidence to suggest that acupuncture may be useful for joint osteoarthritis (OA), mainly due to its ability to stimulate vasodilation and facilitate neovascularization (Ahsin et al., 2009; Li et al., 2009; Loiza et al., 2002; Wu et al., 2010). A number of studies have shown improved blood flow to joints secondary to acupuncture may facilitate the recruitment of opioid producing immune cells required to reduce the level of inflammatory cytokines (Ahsin et al., 2009; Huang et al., 2007; Zhang et al., 2014). There is also limited evidence suggesting that acupuncture may stimulate an increase in hyaluronic acid, allowing the synovial fluid to better lubricate the joint (Li et al., 2009; Wu et al., 2010). Given Scully's mechanism of TMD, (Scully, 2008, 2013) it may therefore be particularly advantageous to target traditional acupoints GB2 and SI19, as they are anatomically positioned directly over the TMJ posterior capsule (Deadman et al., 2007). (See figure 1)

PLACE FIGURE 1 HERE

*Figure 1: Recommended needle insertion sites for the conservative treatment of TMD.*

### **Conclusion**

There is limited evidence to support soft tissue release and strengthening exercises for the muscles of mastication in patients with TMD. For the reduction of pain associated with TMD, there is little to no evidence to support the use of electrophysical modalities and splint management. For the reduction of pain and disability in TMD, non-thrust joint mobilization and HVLA thrust manipulation to the TMJ and/or upper cervical articulations, and dry needling or acupuncture to the lateral pterygoid muscle and posterior peri-articular connective tissue may provide the most evidence-based approach for conservatively treating TMD.

**Works Cited**

- Ahsin S, Saleem S, Bhatti AM, Iles RK, Aslam M 2009 Clinical and endocrinological changes after electro-acupuncture treatment in patients with osteoarthritis of the knee. *Pain* 147, 60-66.
- Al-Ani MZ, Davies SJ, Gray RJ, Sloan P, Glenny AM 2004 Stabilisation splint therapy for temporomandibular pain dysfunction syndrome. *Cochrane Database Syst Rev*, CD002778.
- Alves BM, Macedo CR, Januzzi E, Grossmann E, Atallah AN, Peccin S 2013 Mandibular manipulation for the treatment of temporomandibular disorder. *J Craniofac Surg* 24, 488-493.
- Anderson CR, Boeh SD, Morris RL, Sembrowich WL, Panus PC 2003a Quantification of total dexamethasone phosphate delivery by iontophoresis. *Int J Pharm Compd* 7, 155-159.
- Anderson CR, Morris RL, Boeh SD, Panus PC, Sembrowich WL 2003b Effects of iontophoresis current magnitude and duration on dexamethasone deposition and localized drug retention. *Phys Ther* 83, 161-170.
- Ariji Y, Nakayama M, Nishiyama W, Ogi N, Sakuma S, Katsumata A, Kurita K, Ariji E 2015 Potential clinical application of masseter and temporal muscle massage treatment using an oral rehabilitation robot in temporomandibular disorder patients with myofascial pain. *Cranio* 33, 256-262.
- Armijo-Olivo S, Pitance L, Singh V, Neto F, Thie N, Michelotti A 2016 Effectiveness of Manual Therapy and Therapeutic Exercise for Temporomandibular Disorders: Systematic Review and Meta-Analysis. *Phys Ther* 96, 9-25.
- Ayyildiz S, Emir F, Sahin C 2015 Evaluation of Low-Level Laser Therapy in TMD Patients. *Case Rep Dent* 2015, 424213.
- Bae Y, Park Y 2013 The Effect of Relaxation Exercises for the Masticator Muscles on Temporomandibular Joint Dysfunction (TMD). *J Phys Ther Sci* 25, 583-586.
- Bialosky JE, Bishop MD, Price DD, Robinson ME, George SZ 2009 The mechanisms of manual therapy in the treatment of musculoskeletal pain: a comprehensive model. *Man Ther* 14, 531-538.
- Braun BL 1987 Treatment of an acute anterior disk displacement in the temporomandibular joint. A case report. *Phys Ther* 67, 1234-1236.
- Bronfort G, Haas M, Evans R, Leininger B, Triano J 2010 Effectiveness of manual therapies: the UK evidence report. *Chiropr Osteopat* 18, 3.
- Butts R, Dunning J, Perreault T, Maurad F, Grubb M 2016 Peripheral and Spinal Mechanisms of Pain and Dry Needling Mediated Analgesia: A Clinical Resource Guide for Health Care Professionals. *International Journal of Physical Medicine and Rehabilitation* 216.

- Calixtre LB, Moreira RF, Franchini GH, Alburquerque-Sendin F, Oliveira AB 2015 Manual therapy for the management of pain and limited range of motion in subjects with signs and symptoms of temporomandibular disorder: a systematic review of randomised controlled trials. *J Oral Rehabil* 42, 847-861.
- Cameron M 2003 *Physical Agents in Rehabilitation from Research to Practice: Second Edition*. Saunders, St. Louis, MO.
- Carmeli E, Sheklow S, Bloomenfeld I 2001 Comparative Study of Repositioning Splint Therapy and Passive Manual Range of Motion Techniques for Anterior Displaced Temporomandibular Discs with Unstable Excursive Reduction. *Physiotherapy* 87, 26-36.
- Castroflorio T, Falla D, Wang K, Svensson P, Farina D 2012 Effect of experimental jaw-muscle pain on the spatial distribution of surface EMG activity of the human masseter muscle during tooth clenching. *J Oral Rehabil* 39, 81-92.
- Cho SH, Whang WW 2010 Acupuncture for temporomandibular disorders: a systematic review. *J Orofac Pain* 24, 152-162.
- Choi YJ, Lee JE, Moon WK, Cho SH 2013 Does the effect of acupuncture depend on needling sensation and manipulation? *Complement Ther Med* 21, 207-214.
- Chua NH, Suijlekom HV, Wilder-Smith OH, Vissers KC 2012 Understanding cervicogenic headache. *Anesth Pain Med* 2, 3-4.
- Colloca CJ, Keller TS, Gunzburg R 2004 Biomechanical and neurophysiological responses to spinal manipulation in patients with lumbar radiculopathy. *J Manipulative Physiol Ther* 27, 1-15.
- Deadman P, Al-Khafaji M, Baker KG 2007 *A Manual of Acupuncture 2nd Edition*. Journal of Chinese Medicine Publications, East Sussex, England.
- Desmons S, Graux F, Atassi M, Libersa P, Dupas PH 2007 The lateral pterygoid muscle, a heterogeneous unit implicated in temporomandibular disorder: a literature review. *Cranio* 25, 283-291.
- Dincer F, Linde K 2003 Sham interventions in randomized clinical trials of acupuncture--a review. *Complement Ther Med* 11, 235-242.
- Diracoglu D, Vural M, Karan A, Aksoy C 2012 Effectiveness of dry needling for the treatment of temporomandibular myofascial pain: a double-blind, randomized, placebo controlled study. *J Back Musculoskelet Rehabil* 25, 285-290.
- Dunning J, Butts R, Mourad F, Young I, Flannagan S, Perreault T 2014 Dry needling: a literature review with implications for clinical practice guidelines. *Phys Ther Rev* 19, 252-265.
- Dunning JR, Cleland JA, Waldrop MA, Arnot CF, Young IA, Turner M, Sigurdsson G 2012 Upper cervical and upper thoracic thrust manipulation versus nonthrust mobilization in patients with mechanical neck pain: a multicenter randomized clinical trial. *J Orthop Sports Phys Ther* 42, 5-18.
- Fernandez-Carnero J, La Touche R, Ortega-Santiago R, Galan-del-Rio F, Pesquera J, Ge HY, Fernandez-de-Las-Penas C 2010 Short-term effects of dry needling of active myofascial trigger points in the masseter muscle in patients with temporomandibular disorders. *J Orofac Pain* 24, 106-112.
- Fernandez-de-Las-Penas C, Galan-Del-Rio F, Alonso-Blanco C, Jimenez-Garcia R, Arendt-Nielsen L, Svensson P 2010 Referred pain from muscle trigger points in the



- masticatory and neck-shoulder musculature in women with temporomandibular disorders. *J Pain* 11, 1295-1304.
- Fischer AA 1988 Documentation of myofascial trigger points. *Arch Phys Med Rehabil* 69, 286-291.
- Friedman MH 1997 The hypomobile temporomandibular joint. *Gen Dent* 45, 282-285.
- Fuglevand AJ, Winter DA, Patla AE, Stashuk D 1992 Detection of motor unit action potentials with surface electrodes: influence of electrode size and spacing. *Biol Cybern* 67, 143-153.
- Furto ES, Cleland JA, Whitman JM, Olson KA 2006 Manual physical therapy interventions and exercise for patients with temporomandibular disorders. *Cranio* 24, 283-291.
- Gauer RL, Semidey MJ 2015 Diagnosis and treatment of temporomandibular disorders. *Am Fam Physician* 91, 378-386.
- Goldman N, Chen M, Fujita T, Xu Q, Peng W, Liu W, Jensen TK, Pei Y, Wang F, Han X, Chen JF, Schnermann J, Takano T, Bekar L, Tieu K, Nedergaard M 2010 Adenosine A1 receptors mediate local anti-nociceptive effects of acupuncture. *Nat Neurosci* 13, 883-888.
- Gonzalez-Perez LM, Infante-Cossio P, Granados-Nunez M, Urresti-Lopez FJ 2012 Treatment of temporomandibular myofascial pain with deep dry needling. *Med Oral Patol Oral Cir Bucal* 17, e781-785.
- Gonzalez-Perez LM, Infante-Cossio P, Granados-Nunez M, Urresti-Lopez FJ, Lopez-Martos R, Ruiz-Canela-Mendez P 2015 Deep dry needling of trigger points located in the lateral pterygoid muscle: Efficacy and safety of treatment for management of myofascial pain and temporomandibular dysfunction. *Med Oral Patol Oral Cir Bucal* 20, e326-333.
- Gotou M, Nagata K, Sugawara Y 2010 Short-term effectiveness of the Jog-manipulation technique for temporomandibular disorder (TMD) patients with limited mouth-opening -A randomized controlled trial- *Journal of the Japanese Society for the Temporomandibular Joint* 22, 84-91.
- Grillo CM, Canales Gde L, Wada RS, Alves MC, Barbosa CM, Berzin F, de Sousa Mda L 2015 Could Acupuncture Be Useful in the Treatment of Temporomandibular Dysfunction? *J Acupunct Meridian Stud* 8, 192-199.
- Guarda-Nardini L, Cadorin C, Frizziero A, Masiero S, Manfredini D 2016 Interrelationship between temporomandibular joint osteoarthritis (OA) and cervical spine pain: Effects of intra-articular injection with hyaluronic acid. *Cranio*, 1-7.
- Guarda-Nardini L, Stecco A, Stecco C, Masiero S, Manfredini D 2012 Myofascial pain of the jaw muscles: comparison of short-term effectiveness of botulinum toxin injections and fascial manipulation technique. *Cranio* 30, 95-102.
- Hiraba K, Hibino K, Hiranuma K, Negoro T 2000 EMG activities of two heads of the human lateral pterygoid muscle in relation to mandibular condyle movement and biting force. *J Neurophysiol* 83, 2120-2137.
- Huang J, Zhuo LS, Wang YY, Peng ZL, Huang YR, Wang Y, Yang L 2007 [Effects of electroacupuncture on synovial IL-1beta and TNF-alpha contents in the rabbit with knee osteoarthritis]. *Zhen Ci Yan Jiu* 32, 115-118.

- Jayaseelan DJ, Tow NS 2016 Cervicothoracic junction thrust manipulation in the multimodal management of a patient with temporomandibular disorder. *J Man Manip Ther* 24, 90-97.
- Johnstone DR, Templeton M 1980 The feasibility of palpating the lateral pterygoid muscle. *J Prosthet Dent* 44, 318-323.
- Jung A, Shin BC, Lee MS, Sim H, Ernst E 2011 Acupuncture for treating temporomandibular joint disorders: a systematic review and meta-analysis of randomized, sham-controlled trials. *J Dent* 39, 341-350.
- Kong J, Gollub R, Huang T, Polich G, Napadow V, Hui K, Vangel M, Rosen B, Kaptchuk TJ 2007 Acupuncture de qi, from qualitative history to quantitative measurement. *J Altern Complement Med* 13, 1059-1070.
- Kraus S 1988 Management of the Craniomandibular Complex. Churchill Livingstone, New York.
- Kraus S 2004 Temporomandibular Disorders In: Saunders HD Evaluation, treatment and prevention of musculoskeletal disorders. The Saunders Group, Chaska, MN.
- Kropmans TJ, Dijkstra PU, Stegenga B, Stewart R, de Bont LG 1999a Smallest detectable difference in outcome variables related to painful restriction of the temporomandibular joint. *J Dent Res* 78, 784-789.
- Kropmans TJ, Dijkstra PU, van Veen A, Stegenga B, de Bont LG 1999b The smallest detectable difference of mandibular function impairment in patients with a painfully restricted temporomandibular joint. *J Dent Res* 78, 1445-1449.
- La Touche R, Goddard G, De-la-Hoz JL, Wang K, Paris-Aleman A, Angulo-Diaz-Parreno S, Mesa J, Hernandez M 2010 Acupuncture in the treatment of pain in temporomandibular disorders: a systematic review and meta-analysis of randomized controlled trials. *Clin J Pain* 26, 541-550.
- Lafreniere CM, Lamontagne M, el-Sawy R 1997 The role of the lateral pterygoid muscles in TMJ disorders during static conditions. *Cranio* 15, 38-52.
- Langevin HM, Bouffard NA, Badger GJ, Churchill DL, Howe AK 2006 Subcutaneous tissue fibroblast cytoskeletal remodeling induced by acupuncture: evidence for a mechanotransduction-based mechanism. *J Cell Physiol* 207, 767-774.
- Langevin HM, Bouffard NA, Badger GJ, Iatridis JC, Howe AK 2005 Dynamic fibroblast cytoskeletal response to subcutaneous tissue stretch ex vivo and in vivo. *Am J Physiol Cell Physiol* 288, C747-756.
- Langevin HM, Bouffard NA, Fox JR, Palmer BM, Wu J, Iatridis JC, Barnes WD, Badger GJ, Howe AK 2011 Fibroblast cytoskeletal remodeling contributes to connective tissue tension. *J Cell Physiol* 226, 1166-1175.
- Lauriti L, Motta LJ, de Godoy CH, Biasotto-Gonzalez DA, Politti F, Mesquita-Ferrari RA, Fernandes KP, Bussadori SK 2014 Influence of temporomandibular disorder on temporal and masseter muscles and occlusal contacts in adolescents: an electromyographic study. *BMC Musculoskelet Disord* 15, 123.
- Li ZD, Cao LH, Wang SC 2009 [Effect of moxibustion in treating knee joint osteoarthritis and its relation with contents of hyaluronic acid in serum and synovial fluid]. *Zhongguo Zhong Xi Yi Jie He Za Zhi* 29, 883-885.
- Lin CS 2014 Brain signature of chronic orofacial pain: a systematic review and meta-analysis on neuroimaging research of trigeminal neuropathic pain and temporomandibular joint disorders. *PLoS One* 9, e94300.

- List T, Axelsson S 2010 Management of TMD: evidence from systematic reviews and meta-analyses. *J Oral Rehabil* 37, 430-451.
- Liu ZJ, Wang HY, Pu WY 1989 A comparative electromyographic study of the lateral pterygoid muscle and arthrography in patients with temporomandibular joint disturbance syndrome sounds. *J Prosthet Dent* 62, 229-233.
- Loaiza LA, Yamaguchi S, Ito M, Ohshima N 2002 Electro-acupuncture stimulation to muscle afferents in anesthetized rats modulates the blood flow to the knee joint through autonomic reflexes and nitric oxide. *Auton Neurosci* 97, 103-109.
- Lundeberg T, Lund I 2007 Are reviews based on sham acupuncture procedures in fibromyalgia syndrome (FMS) valid? *Acupunct Med* 25, 100-106.
- Mahan PE, Wilkinson TM, Gibbs CH, Mauderli A, Brannon LS 1983 Superior and inferior bellies of the lateral pterygoid muscle EMG activity at basic jaw positions. *J Prosthet Dent* 50, 710-718.
- Manfredini D, Cocilovo F, Stellini E, Favero L, Guarda-Nardini L 2013 Surface electromyography findings in unilateral myofascial pain patients: comparison of painful vs. non painful sides. *Pain Med* 14, 1848-1853.
- Mansilla-Ferragut P, Fernandez-de-Las Penas C, Alburquerque-Sendin F, Cleland JA, Bosca-Gandia JJ 2009 Immediate effects of atlanto-occipital joint manipulation on active mouth opening and pressure pain sensitivity in women with mechanical neck pain. *J Manipulative Physiol Ther* 32, 101-106.
- Mapelli A, Zanandrea Machado BC, Giglio LD, Sforza C, De Felicio CM 2016 Reorganization of muscle activity in patients with chronic temporomandibular disorders. *Arch Oral Biol* 72, 164-171.
- Martins WR, Blasczyk JC, Aparecida Furlan de Oliveira M, Lagoa Goncalves KF, Bonini-Rocha AC, Dugailly PM, de Oliveira RJ 2016 Efficacy of musculoskeletal manual approach in the treatment of temporomandibular joint disorder: A systematic review with meta-analysis. *Man Ther* 21, 10-17.
- McMillan AS, Nolan A, Kelly PJ 1997 The efficacy of dry needling and procaine in the treatment of myofascial pain in the jaw muscles. *J Orofac Pain* 11, 307-314.
- McNamara JA, Jr. 1973 The independent functions of the two heads of the lateral pterygoid muscle. *Am J Anat* 138, 197-205.
- McNeely ML, Armijo Olivo S, Magee DJ 2006 A systematic review of the effectiveness of physical therapy interventions for temporomandibular disorders. *Phys Ther* 86, 710-725.
- Mesa-Jimenez JA, Sanchez-Gutierrez J, de-la-Hoz-Aizpurua JL, Fernandez-de-las-Penas C 2015 Cadaveric validation of dry needle placement in the lateral pterygoid muscle. *J Manipulative Physiol Ther* 38, 145-150.
- Miernik M, Wieckiewicz M, Paradowska A, Wieckiewicz W 2012 Massage therapy in myofascial TMD pain management. *Adv Clin Exp Med* 21, 681-685.
- Mina R, Melson P, Powell S, Rao M, Hinze C, Passo M, Graham TB, Brunner HI 2011 Effectiveness of dexamethasone iontophoresis for temporomandibular joint involvement in juvenile idiopathic arthritis. *Arthritis Care Res (Hoboken)* 63, 1511-1516.
- Moller E, Sheikholeslam A, Lous I 1984 Response of elevator activity during mastication to treatment of functional disorders. *Scand J Dent Res* 92, 64-83.

- Monaco A, Cozzolino V, Cattaneo R, Cutilli T, Spadaro A 2008 Osteopathic manipulative treatment (OMT) effects on mandibular kinetics: kinesiographic study. *Eur J Paediatr Dent* 9, 37-42.
- Moore K, Dalley A 2006 *Clinically Oriented Anatomy: Fifth Edition*. Lippincott Williams & Wilkins, Philadelphia.
- Moraes A 2013 Therapeutic exercises for the control of temporomandibular disorders. *Dental Press J Orthod* 18, 134-139.
- Mulet M, Decker KL, Look JO, Lenton PA, Schiffman EL 2007 A randomized clinical trial assessing the efficacy of adding 6 x 6 exercises to self-care for the treatment of masticatory myofascial pain. *J Orofac Pain* 21, 318-328.
- Murray GM, Peck CC 2007 Orofacial pain and jaw muscle activity: a new model. *J Orofac Pain* 21, 263-278; discussion 279-288.
- Murray GM, Phanachet I, Uchida S, Whittle T 2004 The human lateral pterygoid muscle: a review of some experimental aspects and possible clinical relevance. *Aust Dent J* 49, 2-8.
- Nagata K, Maruyama H, Mizuhashi R, Morita S, Hori S, Yokoe T, Sugawara Y 2015 Efficacy of stabilisation splint therapy combined with non-splint multimodal therapy for treating RDC/TMD axis I patients: a randomised controlled trial. *J Oral Rehabil*.
- Ogawa M, Kanbe T, Kano A, Kubota F, Makiguchi T, Miyazaki H, Yokoo S 2015 Conservative reduction by lever action of chronic bilateral mandibular condyle dislocation. *Cranio* 33, 142-147.
- Oliveira-Campelo NM, Rubens-Rebelatto J, Marti NVFJ, Alburquerque-Sendi NF, Fernandez-de-Las-Penas C 2010 The immediate effects of atlanto-occipital joint manipulation and suboccipital muscle inhibition technique on active mouth opening and pressure pain sensitivity over latent myofascial trigger points in the masticatory muscles. *J Orthop Sports Phys Ther* 40, 310-317.
- Peck CC, Murray GM, Gerzina TM 2008 How does pain affect jaw muscle activity? The Integrated Pain Adaptation Model. *Aust Dent J* 53, 201-207.
- Pihut M, Ferendiuk E, Szewczyk M, Kasprzyk K, Wieckiewicz M 2016 The efficiency of botulinum toxin type A for the treatment of masseter muscle pain in patients with temporomandibular joint dysfunction and tension-type headache. *J Headache Pain* 17, 29.
- Pinho JC, Caldas FM, Mora MJ, Santana-Penin U 2000 Electromyographic activity in patients with temporomandibular disorders. *J Oral Rehabil* 27, 985-990.
- Porporatti A, Costa Y, Stuginski-Borbosa J, Bonjardim L, Conti P 2015 Acupuncture therapeutic protocols for the management of temporomandibular disorders. *Sao Paulo* 16.
- Qvintus V, Suominen AL, Huttunen J, Raustia A, Ylostalo P, Sipila K 2015 Efficacy of stabilisation splint treatment on facial pain - 1-year follow-up. *J Oral Rehabil* 42, 439-446.
- Reid KI, Dionne RA, Sicard-Rosenbaum L, Lord D, Dubner RA 1994 Evaluation of iontophoretically applied dexamethasone for painful pathologic temporomandibular joints. *Oral Surg Oral Med Oral Pathol* 77, 605-609.
- Robertson VJ, Baker KG 2001 A review of therapeutic ultrasound: effectiveness studies. *Phys Ther* 81, 1339-1350.

- Rocabado M, Iglarsh Z 1991 Musculoskeletal approach to maxillofacial pain. JB Lippincott, Philadelphia, PA.
- Rubis LM, Rubis D, Winchester B 2014 A collaborative approach between chiropractic and dentistry to address temporomandibular dysfunction: a case report. *J Chiropr Med* 13, 55-61.
- Saghafi D, Curl DD 1995 Chiropractic manipulation of anteriorly displaced temporomandibular disc with adhesion. *J Manipulative Physiol Ther* 18, 98-104.
- Sandberg M, Lundeberg T, Lindberg LG, Gerdle B 2003 Effects of acupuncture on skin and muscle blood flow in healthy subjects. *Eur J Appl Physiol* 90, 114-119.
- Schiffman EL, Braun BL, Lindgren BR 1996 Temporomandibular joint iontophoresis: a double-blind randomized clinical trial. *J Orofac Pain* 10, 157-165.
- Sciotti VM, Mittak VL, DiMarco L, Ford LM, Plezbert J, Santipadri E, Wigglesworth J, Ball K 2001 Clinical precision of myofascial trigger point location in the trapezius muscle. *Pain* 93, 259-266.
- Scully C 2008 *Oral and Maxillofacial Medicine: the Basis of Diagnosis and Treatment* (2nd ed). Churchill Livingstone, Edinburgh.
- Scully C 2013 *Oral and Maxillofacial Medicine: the Basis of Diagnosis and Treatment* (3rd ed). Churchill Livingstone, Edinburgh.
- Shaffer SM, Brismee JM, Sizer PS, Courtney CA 2014 Temporomandibular disorders. Part 2: conservative management. *J Man Manip Ther* 22, 13-23.
- Shanks P, Curran M, Fletcher P, Thompson R 2010 The effectiveness of therapeutic ultrasound for musculoskeletal conditions of the lower limb: A literature review. *Foot (Edinb)* 20, 133-139.
- Shimada A, Baad-Hansen L, Svensson P 2015 Effect of experimental jaw muscle pain on dynamic bite force during mastication. *Arch Oral Biol* 60, 256-266.
- Simons D, Travell J, Simons L 1999 *Travell & Simons' myofascial pain and dysfunction : the trigger point manual*. Lippincott Williams & Wilkins, Baltimore.
- Simons DG, Travell JG 1983 Myofascial origins of low back pain. 1. Principles of diagnosis and treatment. *Postgrad Med* 73, 66, 68-70, 73 passim.
- Smutok MA, Mayo MF, Gabaree CL, Ferslew KE, Panus PC 2002 Failure to detect dexamethasone phosphate in the local venous blood postcathodic iontophoresis in humans. *J Orthop Sports Phys Ther* 32, 461-468.
- Spadaro A, Ciarrocchi I, Masci C, Cozzolino V, Monaco A 2014 Effects of intervertebral disc disorders of low back on the mandibular kinematic: kinesiographic study. *BMC Res Notes* 7, 569.
- Stelzenmueller W, Umstadt H, Weber D, Goenner-Oezkan V, Kopp S, Lisson J 2016 Evidence - The intraoral palpability of the lateral pterygoid muscle - A prospective study. *Ann Anat* 206, 89-95.
- Takano T, Chen X, Luo F, Fujita T, Ren Z, Goldman N, Zhao Y, Markman JD, Nedergaard M 2012 Traditional acupuncture triggers a local increase in adenosine in human subjects. *J Pain* 13, 1215-1223.
- Taylor M, Suvinen T, Reade P 1994 The effect of Grade IV distraction mobilization on patients with temporomandibular pain-dysfunction disorder. *Physiother Theory Pract* 10.

- Tosato Jde P, Caria PH, Gomes CA, Berzin F, Politti F, Gonzalez Tde O, Biasotto-Gonzalez DA 2015 Correlation of stress and muscle activity of patients with different degrees of temporomandibular disorder. *J Phys Ther Sci* 27, 1227-1231.
- Tough EA, White AR, Richards S, Campbell J 2007 Variability of criteria used to diagnose myofascial trigger point pain syndrome--evidence from a review of the literature. *Clin J Pain* 23, 278-286.
- Turp JC, Minagi S 2001 Palpation of the lateral pterygoid region in TMD--where is the evidence? *J Dent* 29, 475-483.
- van der Windt DA, van der Heijden GJ, van den Berg SG, ter Riet G, de Winter AF, Bouter LM 1999 Ultrasound therapy for musculoskeletal disorders: a systematic review. *Pain* 81, 257-271.
- White A, Cummings M, Barlas P, Cardini F, Filshie J, Foster NE, Lundeberg T, Stener-Victorin E, Witt C 2008 Defining an adequate dose of acupuncture using a neurophysiological approach--a narrative review of the literature. *Acupunct Med* 26, 111-120.
- White A, Foster NE, Cummings M, Barlas P 2007 Acupuncture treatment for chronic knee pain: a systematic review. *Rheumatology (Oxford)* 46, 384-390.
- Wieckiewicz M, Boening K, Wiland P, Shiau YY, Paradowska-Stolarz A 2015 Reported concepts for the treatment modalities and pain management of temporomandibular disorders. *J Headache Pain* 16, 106.
- Wu MX, Li XH, Lin MN, Jia XR, Mu R, Wan WR, Chen RH, Chen LH, Lin WQ, Huang CY, Zhang XR, Hong KD, Li L, Liu XX 2010 Clinical study on the treatment of knee osteoarthritis of Shen-Sui insufficiency syndrome type by electroacupuncture. *Chin J Integr Med* 16, 291-297.
- Yabe T, Tsuda T, Hirose S, Ozawa T, Kawai K 2014 Treatment of acute temporomandibular joint dislocation using manipulation technique for disk displacement. *J Craniofac Surg* 25, 596-597.
- Yim YK, Lee H, Hong KE, Kim YI, Lee BR, Son CG, Kim JE 2007 Electro-acupuncture at acupoint ST36 reduces inflammation and regulates immune activity in Collagen-Induced Arthritic Mice. *Evid Based Complement Alternat Med* 4, 51-57.
- Zhang R, Lao L, Ren K, Berman BM 2014 Mechanisms of acupuncture-electroacupuncture on persistent pain. *Anesthesiology* 120, 482-503.
- Zhou W, Benharash P 2014 Significance of "Deqi" response in acupuncture treatment: myth or reality. *J Acupunct Meridian Stud* 7, 186-189.



| Acupoints | Associated Anatomy   |
|-----------|--|
| ST5       | Anterior aspect of distal masseter muscle  |
| ST6       | Posterior aspect of distal masseter muscle   |
| ST7       | Proximal masseter muscle, superficially<br>Inferior head of lateral pterygoid muscle, deep |
| ST8       | Anterosuperior temporalis muscle   |
| Taiyang   | Anteroinferior temporalis muscle   |
| GB2       | Posterior TMJ capsule  |
| SI19      | Posterior TMJ capsule  |

Figure 1: Recommended needle insertion sites for the conservative treatment of TMD.